

CYCLIC SIEVING PHENOMENON ON ANNULAR NONCROSSING PERMUTATIONS

JANG SOO KIM

ABSTRACT. We show an instance of the cyclic sieving phenomenon on annular noncrossing permutations with given cycle types. We define annular q -Kreweras numbers, annular q -Narayana numbers, and annular q -Catalan number, all of which are polynomials in q . We then show that these polynomials exhibit the cyclic sieving phenomenon on annular noncrossing permutations. We also show that a sum of annular q -Kreweras numbers becomes an annular q -Narayana number and a sum of q -Narayana numbers becomes an annular q -Catalan number.

1. INTRODUCTION

Let π be a permutation of $[n] = \{1, 2, \dots, n\}$. One can represent π inside a disk as shown in Figure 1. If the arrows of the diagram of π are noncrossing and if every cycle of π is oriented clockwise, then π is called a *noncrossing permutation*. If we replace each cycle with a block, then we get a bijection from noncrossing permutations and noncrossing partitions.

It is well known that the number of noncrossing permutations of $[n]$ is the *Catalan number*

$$\text{Cat}(n) = \frac{1}{n+1} \binom{2n}{n},$$

and the number of noncrossing permutations of $[n]$ with k cycles is the *Narayana number*

$$\text{Nara}(n, k) = \frac{1}{n} \binom{n}{k-1} \binom{n}{k}.$$

We denote by $\text{Par}(n, k)$ the set of integer partitions of n with k parts. If λ has m_i parts of size i for $i = 1, 2, \dots$, then we also write as $\lambda = (1^{m_1}, 2^{m_2}, \dots)$. We will use the following notations: if $\lambda = (1^{m_1}, 2^{m_2}, \dots)$ has k parts, i.e. $m_1 + m_2 + \dots = k$, we define

$$\binom{k}{\lambda} = \frac{k!}{m_1! m_2! \dots}, \quad [k]_q = \frac{[k]_q!}{[m_1]_q! [m_2]_q! \dots},$$

Date: October 30, 2012.




FIGURE 1. A representation of the permutation $(1, 3, 4, 5)(2)(6, 10)(7, 8, 9)$ inside a disk.

where we use the standard notations

$$[n]_q = \frac{1 - q^n}{1 - q}, \quad [n]_q! = [1]_q [2]_q \cdots [n]_q, \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n - k]_q!}.$$

The *cycle type* of a permutation π is the partition $\lambda = (1^{m_1}, 2^{m_2}, \dots)$, where m_i is the number of cycles with i elements. Kreweras [7] showed that the number of noncrossing permutations of $[n]$ with cycle type $\lambda = (1^{m_1}, 2^{m_2}, \dots) \in \text{Par}(n, k)$ is equal to the *Kreweras number*

$$\text{Kre}(\lambda) := \frac{1}{k} \binom{n}{k-1} \binom{k}{\lambda}.$$

Bessis and Reiner [2, Theorem 6.2] showed that if X is the set of noncrossing partitions of $[n]$ with type $\lambda = (1^{m_1}, 2^{m_2}, \dots) \in \text{Par}(n, k)$,

$$X(q) = \frac{1}{[k]_q} \begin{bmatrix} n \\ k-1 \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q,$$

and C is the cyclic group of rotations acting on X , then $(X, X(q), C)$ exhibits the cyclic sieving phenomenon, see Section 2 for the definition.

Reiner and Sommers [10] defined the *q-Kreweras number* for $\lambda = (1^{m_1}, 2^{m_2}, \dots) \in \text{Par}(n, k)$ by

$$\text{Kre}_q(\lambda) = \frac{q^{(n+1)(n-k) - \tau(\lambda)}}{[k]_q} \begin{bmatrix} n \\ k-1 \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q,$$

where $\tau(\lambda) = \sum_{i \geq 1} \lambda'_i \lambda'_{i+1}$, and the *q-Narayana number* by

$$\text{Nara}_q(n, k) = \frac{q^{(n-k)(n+1-k)}}{[n]_q} \begin{bmatrix} n \\ k-1 \end{bmatrix}_q \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

Reiner and Sommers [10] showed that

$$\sum_{\lambda \in \text{Par}(n, k)} \text{Kre}_q(\lambda) = \text{Nara}_q(n, k)$$

and

$$\sum_{k=0}^n \text{Nara}_q(n, k) = \text{Cat}_q(n),$$

where $\text{Cat}_q(n)$ is the *q-Catalan number* defined by

$$\text{Cat}_q(n) = \frac{1}{[n+1]_q} \begin{bmatrix} 2n \\ n \end{bmatrix}_q.$$

In this paper we prove analogous results for annular noncrossing permutations. Annular noncrossing permutations are an annulus-analog of noncrossing permutations. They were first introduced by Mingo and Nica [8] and studied further in [4, 9]. Recently Kim, Seo, and Shin [6] used annular noncrossing permutations to give a combinatorial proof of Goulden and Jackson's formula [3] for the number of minimal transitive factorizations of a product of two cycles.

This paper is organized as follows. In Section 2 we define annular noncrossing permutations and give an instance of the cyclic sieving phenomenon on them. In Section 3 we define annular *q-Kreweras numbers*, three types of annular *q-Narayana numbers*, and annular *q-Catalan numbers* and show that a sum of annular *q-Kreweras numbers* becomes an annular *q-Narayana number*, and a sum of annular *q-Narayana numbers* becomes an annular *q-Catalan number*. We then show that these polynomials give cyclic sieving phenomena. In Section 4 we enumerate annular noncrossing matchings.




FIGURE 2. A representation of the annular noncrossing permutation $(1, 2, 3, 6, 15, 10, 11)(4, 5)(7, 8, 9, 13, 14)(12)$.

2. ENUMERATION OF ANNULAR NONCROSSING PERMUTATIONS

Let n and m be positive integers. An (n, m) -annulus is an annulus in which $1, 2, \dots, n$ are arranged in clockwise order on the exterior circle and $n + 1, n + 2, \dots, n + m$ are arranged in counter-clockwise order on the interior circle.

Let $(a_1, \dots, a_k)$ be a cycle whose elements are contained in $[n + m]$. We will represent this cycle inside an (n, m) -annulus by drawing an arrow from a_i to a_{i+1} for each $i = 1, 2, \dots, k$, where $a_{k+1} = a_1$. An *interior cycle* (respectively *exterior cycle*) is a cycle all of whose elements are on the interior (respectively exterior) circle. A *connected cycle* is a cycle which contains both an element in the interior circle and an element in the exterior cycle. Suppose $\{e_1 < e_2 < \dots < e_u\} = \{a_1, \dots, a_k\} \cap [n]$ and $\{i_1 < i_2 < \dots < i_v\} = \{a_1, \dots, a_k\} \cap \{n + 1, n + 2, \dots, n + m\}$. Then we say that the cycle $(a_1, \dots, a_k)$ is *oriented clockwise* if we can express $(a_1, \dots, a_k) = (e'_1, \dots, e'_u, i'_1, \dots, i'_v)$ for some integers $e'_1, \dots, e'_u, i'_1, \dots, i'_v$ with $(e'_1, \dots, e'_u) = (e_1, \dots, e_u)$ and $(i'_1, \dots, i'_v) = (i_1, \dots, i_v)$. In this case we say that the cycle $(a_1, \dots, a_k)$ is of *size* k , of *exterior size* u and of *interior size* v .

A permutation of $[n + m]$ is called an (n, m) -annular noncrossing permutation if we can draw its cycles inside an (n, m) -annulus in such a way that every cycle is oriented clockwise and there are no crossing arrows, see Figure 2.

Remark 1. Unlike noncrossing permutations, the map changing each cycle to a block is not a one-to-one correspondence from annular noncrossing permutations to annular noncrossing partitions. However, as is shown in [8, Proposition 4.4], if there are at least two connected cycles, then this map becomes a bijection. Thus every result in this paper on annular noncrossing permutations with at least two connected cycles works for annular noncrossing partitions.

If an (n, m) -annular noncrossing permutation has a connected cycle, it is called *connected*. Since a disconnected annular noncrossing permutation is essentially a disjoint union of two noncrossing permutations, in this paper we will only consider connected annular noncrossing permutations.

We denote by $\text{ANC}(n, m)$ the set of connected (n, m) -annular noncrossing permutations. For $\pi \in \text{ANC}(n, m)$, the *exterior cycle type* (respectively *interior cycle type*) of π is the partition $(1^{m_1}, 2^{m_2}, \dots)$ where m_i is the number of exterior cycles (respectively interior cycles) of size i . The *connected exterior cycle type* (respectively *connected interior cycle type*) of π is the partition $(1^{m_1}, 2^{m_2}, \dots)$ where m_i is the number of connected cycles of exterior size (respectively interior size) i .

For integers $n, m, c, r, s, R, S \geq 0$ and $\alpha \in \text{Par}(R, r)$, $\beta \in \text{Par}(S, s)$, $\lambda \in \text{Par}(n - R, c)$, and $\mu \in \text{Par}(m - S, c)$, we define the following:

- $\text{ANC}(n, m; c)$ is the set of $\pi \in \text{ANC}(n, m)$ with c connected cycles.
- $\text{ANC}(n, m; c, r, s)$ is the set of $\pi \in \text{ANC}(n, m; c)$ with r exterior cycles and s interior cycles.

- $\text{ANC}(n, m; c, r, s, R, S)$ is the set of $\pi \in \text{ANC}(n, m; c, r, s)$ such that the total size of exterior cycles is R and the total size of interior cycles is S .
- $\text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)$ is the set of $\pi \in \text{ANC}(n, m)$ with exterior cycle type $\alpha \in \text{Par}(R, r)$, interior cycle type $\beta \in \text{Par}(S, s)$, connected exterior cycle type $\lambda \in \text{Par}(n-R, c)$, and connected interior cycle type $\mu \in \text{Par}(m-S, c)$.

Definition 1. Suppose a cyclic group C of order n acts on a finite set X . Let $X(q)$ be a polynomial in q . We say that $(X, X(q), C)$ exhibits the *cyclic sieving phenomenon* (CSP) if $X(w(c)) = |\{x \in X : c(x) = x\}|$ for all $c \in C$. Here, $w : C \rightarrow \mathbb{C}^\times$ is an embedding of C into the multiplicative group $\mathbb{C}^\times$ of nonzero complex numbers sending a cyclic generator of C to a primitive n th root of unity.

The CSP was first introduced by Reiner, Stanton, and White [11]. Recently many instances of the CSP have been found. In [12], Sagan gives a nice survey on the CSP.

Usually the CSP is defined with a polynomial $X(q)$ whose coefficients are integers. In this paper we do not assume any conditions on the coefficients of $X(q)$. In fact we will see an instance of the CSP with $X(q)$ whose coefficients are not integers. However, by the definition of the CSP, $X(w(c))$ is always a nonnegative integer.

The goal of this section is to find an instance of the CSP for $\text{ANC}(n, m)$. To this end let $C_1 \times C_2$ be the product of two cyclic groups C_1 acting on the exterior circle and C_2 acting on the interior circle. Then $C_1 \times C_2$ gives a bicyclic action on $\text{ANC}(n, m)$ by $(c_1, c_2)\pi = c_1(c_2(\pi)) = c_2(c_1(\pi))$. One may wonder if this bicyclic action gives a “bicyclic sieving phenomenon” as in [1]. However, this is not the case by the next proposition.

Proposition 2.1. *Let $(c_1, c_2) \in C_1 \times C_2$. Then (c_1, c_2) has no fixed points in $\text{ANC}(n, m)$ unless c_1 and c_2 have the same order.*

Proof. Let d_1 and d_2 be the orders of c_1 and c_2 respectively. Suppose $\pi \in \text{ANC}(n, m)$ is a fixed point, i.e. $(c_1, c_2)\pi = \pi$. Let $k_1 = n/d_1$ and $k_2 = m/d_2$. We can assume that c_1 is the map sending $i \in [n]$ to $j \in [n]$ with $j \equiv i + k_1 \pmod{n}$, and c_2 is the map sending $n + i \in \{n+1, \dots, n+m\}$ to $n + j \in \{n+1, \dots, n+m\}$ with $j \equiv i - k_2 \pmod{m}$. Note that for each $i \in [n]$ (respectively $i \in \{n+1, \dots, n+m\}$) $c_1^{(t)}(i) = i$ (respectively $c_2^{(t)}(i) = i$) implies that t divides d_1 (respectively d_2). Consider a connected cycle $\gamma = (a_1, \dots, a_u, b_1, \dots, b_v)$ of π where $a_1, \dots, a_u \in [n]$ and $b_1, \dots, b_v \in \{n+1, \dots, n+m\}$. Note that $u > 0$ and $v > 0$. Since $(c_1, c_2)^{(d_1)}(\pi) = \pi$ and $(c_1, c_2)^{(d_1)}(\gamma) = (a_1, \dots, a_u, c_2^{(d_1)}(b_1), \dots, c_2^{(d_1)}(b_v))$, we have $(c_1, c_2)^{(d_1)}(\gamma) = \gamma$. In particular, $c_2^{(d_1)}(b_1) = b_1$, which implies that d_1 divides d_2 . Similarly we get that d_2 divides d_1 . Thus $d_1 = d_2$. $\square$

Thus we will only consider the elements (c_1, c_2) for which c_1 and c_2 have the same order. We call such pair (c_1, c_2) an (n, m) -annular rotation, or simply an *annular rotation*. Note that if (c_1, c_2) is an (n, m) -annular rotation, then the order d of this action divides both n and m .

For a partition $\lambda = (1^{m_1}, 2^{m_2}, \dots)$ we denote by $\mathfrak{S}(\lambda)$ the set of rearrangements of the following sequence

$$\underbrace{1, \dots, 1}_{m_1}, \underbrace{2, \dots, 2}_{m_2}, \dots$$

In other words, each element in $\mathfrak{S}(\lambda)$ is a sequence $(a_1, a_2, \dots)$ where each integer i appears exactly m_i times. If each m_i is divisible by d , we define $\lambda/d = (1^{m_1/d}, 2^{m_2/d}, \dots)$. In this case we say that λ is *divisible* by d .

Theorem 2.2. *The number of $\pi \in \text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)$ invariant under an annular rotation of order d is equal to*

$$d \cdot \frac{(\widehat{n} - \widehat{R})(\widehat{m} - \widehat{S})}{\widehat{c}} \binom{\widehat{n}}{\widehat{r}} \binom{\widehat{m}}{\widehat{s}} \binom{\widehat{r}}{\widehat{\alpha}} \binom{\widehat{s}}{\widehat{\beta}} \binom{\widehat{c}}{\widehat{\lambda}} \binom{\widehat{c}}{\widehat{\mu}},$$

if all of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ are divisible by d , and 0 otherwise. Here $\widehat{Z}$ means Z/d .

Sketch of Proof. Since the proof is similar to those in [4, Proposition 4.2] and in [5, Proposition 4.1], we will only give a sketch.

If $d = 1$, this follows from Theorem 2.2. Suppose $d > 1$. It is not difficult to see that if an annular noncrossing permutation π is invariant under an annular rotation of order d , then π has no cycles invariant under this rotation. Thus if any of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ is not divisible by d , then there is no such π . So we will assume that all of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ are divisible by d .

Let (c_1, c_2) be an annular rotation of order d . We will find a bijection between the set

$$\mathcal{A} = \{(\gamma, \pi) : \pi \in \text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu), (c_1, c_2)\pi = \pi, \gamma \text{ is a connected cycle of } \pi\}$$

and the set

$$\mathcal{B} = \left\{ (a, b, R^E, R^I, V^E, V^I, V^{CE}, V^{CI}) \left| \begin{array}{l} a \in [n - R], b \in [m - S], \\ R^E \subset [\hat{n}], R^I \subset [\hat{m}], |R^E| = \hat{r}, |R^I| = \hat{s}, \\ V^E \in \mathfrak{S}(\hat{\alpha}), V^I \in \mathfrak{S}(\hat{\beta}), V^{CE} \in \mathfrak{S}(\hat{\lambda}), V^{CI} \in \mathfrak{S}(\hat{\mu}) \end{array} \right. \right\}.$$

Suppose $(\gamma, \pi) \in \mathcal{A}$. Let $\gamma = (a_1, \dots, a_u, b_1, \dots, b_v)$ with $a_1, \dots, a_u \in [n]$ and $b_1, \dots, b_v \in [n + 1, \dots, n + m]$. We define $a = a_1$ and $b = b_v$. Arrange the integers in $[n + m]$ as follows:

$$(1) \quad a, a + 1, \dots, n, 1, 2, \dots, a - 1, b + 1, b + 2, \dots, n + m, n + 1, n + 2, \dots, b.$$

For each exterior or interior cycle of size t , we place a right parenthesis $)_t$ labeled by t after the rightmost integer in the sequence (1) which is an element of the cycle. We define R^E (respectively R^I) to be the set of integers i with a right parenthesis such that $i \in [\hat{n}]$ (respectively $i - n \in [\hat{m}]$). Then we have $|R^E| = \hat{r}$ and $|R^I| = \hat{s}$. Let $i_1 < i_2 < \dots < i_{\hat{r}}$ be the elements of R^E . We define V^E to be the sequence $(\ell_1, \ell_2, \dots, \ell_{\hat{r}})$ where ℓ_j is the label of the right parenthesis after i_j . The sequence V^I is defined similarly.

Now remove the integers contained in an exterior or an interior cycle from the sequence (1). For each connected cycle, we place a left (respectively right) parenthesis before (respectively after) the leftmost (respectively rightmost) integer in the remaining sequence which is an element of the cycle. Then the c left parentheses divide the first part of the remaining sequence consisting of integers at most n into c subsequences. Let V^{CE} be the sequence of sizes of the c subsequences. Similarly, the c right parentheses divide the second part of the remaining sequence consisting of integers greater than n into c subsequences. We define V^{CI} to be the sequence of sizes of the c subsequences. We have just constructed the map $(\gamma, \pi) \mapsto (a, b, R^E, R^I, V^E, V^I, V^{CE}, V^{CI})$. Using the ideas of [4, Proposition 4.2] and [5, Proposition 4.1], one can show that this gives a bijection from $\mathcal{A}$ to $\mathcal{B}$.

Thus the number of $\pi \in \text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)$ invariant under an annular rotation of order d is $|\mathcal{A}|/c = |\mathcal{B}|/c$, which is easily seen to equal to the number in the theorem. $\square$

Now we state the main theorem of this section.

Theorem 2.3. *Suppose that the cyclic group C of (n, m) -annular rotations acts on the set*

$$X = \text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu).$$

Let

$$X(q) = \frac{[(n - R)(m - S)]_q}{[c]_q} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} r \\ \alpha \end{bmatrix}_q \begin{bmatrix} s \\ \beta \end{bmatrix}_q \begin{bmatrix} c \\ \lambda \end{bmatrix}_q \begin{bmatrix} c \\ \mu \end{bmatrix}_q.$$

Then $(X, X(q), C)$ exhibits the CSP.

We first prove that the $X(q)$ in Theorem 2.3 is a polynomial in q . We need two lemmas.

Lemma 2.4. [11, Proposition 10.1 (iii)] *If $f(q) = h(q)/[k]_q \in \mathbb{Z}[q]$, and $h(q) \in \mathbb{N}[q]$ has symmetric, unimodal coefficient sequence, then $f(q) \in \mathbb{N}[q]$.*

Lemma 2.5. *For nonnegative integers N, n, k with $N \geq n$ and a partition $\lambda \in \text{Par}(n, k)$, we have*

$$\frac{[n]_q}{[k]_q} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q \in \mathbb{N}[q], \quad \frac{[N - n]_q}{[N]_q} \begin{bmatrix} N \\ k \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q \in \mathbb{N}[q].$$

Proof. We claim that

$$(2) \quad \frac{[n]_q}{[k]_q} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q = \frac{1 - q^n}{1 - q^k} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q \in \mathbb{Z}[q].$$

Assuming the claim let us show the lemma. By (2) and the fact $[N]_q = [N - n]_q + q^{N-n}[n]_q$ we also have

$$\begin{aligned} \frac{[N - n]_q}{[N]_q} \begin{bmatrix} N \\ k \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q &= \frac{[N]_q - q^{N-n}[n]_q}{[N]_q} \begin{bmatrix} N \\ k \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q \\ &= \begin{bmatrix} N \\ k \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q - q^{N-n} \begin{bmatrix} N - 1 \\ k - 1 \end{bmatrix}_q \cdot \frac{[n]_q}{[k]_q} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q \in \mathbb{Z}[q]. \end{aligned}$$

Since both $\frac{[n]_q}{[k]_q} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q$ and $\frac{[N - n]_q}{[N]_q} \begin{bmatrix} N \\ k \end{bmatrix}_q \begin{bmatrix} k \\ \lambda \end{bmatrix}_q$ satisfy the conditions in Lemma 2.4 we obtain the lemma.

We now show (2). We will use the q -Pochhammer symbol $(q; q)_r = (1 - q)(1 - q^2) \cdots (1 - q^r)$. Note that $\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$.

Since

$$q^k - 1 = \prod_{j=1}^k (q - \omega_j),$$

where $\omega_1, \omega_2, \dots, \omega_k$ are the k th roots of unity, in order to prove (2) it is sufficient to show that $q - \omega_j$ divides $(1 - q^n) \begin{bmatrix} k \\ \lambda \end{bmatrix}_q$ for all $j = 1, 2, \dots, k$. Fix an integer j and suppose ω_j is a primitive r th root of unity. Then r divides k . Note that $q - \omega_j$ divides $q^s - 1$ if and only if r divides s . Note also that the multiplicity of $q - \omega_j$ as a factor of $q^s - 1$ is at most 1. Thus the multiplicity of the factor $q - \omega_j$ in $(q; q)_s$ is equal to $\lfloor s/r \rfloor$. We have two cases as follows.

CASE 1: r divides n . Then $q - \omega_j$ divides $q^n - 1$ and we are done.

CASE 2: r does not divide n . Let $\lambda = (1^{m_1}, 2^{m_2}, \dots, \ell^{m_\ell})$. Then we have $n = \sum_{i=1}^\ell i \cdot m_i$, $k = \sum_{i=1}^\ell m_i$, and

$$(3) \quad \begin{bmatrix} k \\ \lambda \end{bmatrix}_q = \frac{(q; q)_{m_1+m_2+\dots+m_\ell}}{(q; q)_{m_1} (q; q)_{m_2} \cdots (q; q)_{m_\ell}}.$$

The multiplicities of the factor $q - \omega_j$ in the numerator and denominator of (3) are respectively $\lfloor k/r \rfloor$ and $\lfloor m_1/r \rfloor + \lfloor m_2/r \rfloor + \dots + \lfloor m_\ell/r \rfloor$. Since r does not divide n , at least one of $m_1, m_2, \dots, m_\ell$ is not a multiple of r . Thus we have

$$\left\lfloor \frac{m_1}{r} \right\rfloor + \left\lfloor \frac{m_2}{r} \right\rfloor + \dots + \left\lfloor \frac{m_\ell}{r} \right\rfloor < \frac{m_1}{r} + \frac{m_2}{r} + \dots + \frac{m_\ell}{r} = \frac{k}{r} = \left\lfloor \frac{k}{r} \right\rfloor,$$

which implies that $q - \omega_j$ divides $\begin{bmatrix} k \\ \lambda \end{bmatrix}_q$. This finishes the proof of (2). $\square$

Proposition 2.6. *The $X(q)$ in Theorem 2.3 is a polynomial in q .*

Proof. By Lemma 2.5, $\frac{[m-S]_q}{[c]_q} \begin{bmatrix} c \\ \mu \end{bmatrix}_q$ is a polynomial. Since

$$(4) \quad X(q) = \frac{[(n-R)(m-S)]_q}{[m-S]_q} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} r \\ \alpha \end{bmatrix}_q \begin{bmatrix} s \\ \beta \end{bmatrix}_q \begin{bmatrix} c \\ \lambda \end{bmatrix}_q \frac{[m-S]_q}{[c]_q} \begin{bmatrix} c \\ \mu \end{bmatrix}_q,$$

and $[ab]_q = [a]_q [b]_{q^a} = [b]_q [a]_{q^b}$, we obtain that $X(q)$ is a polynomial in q . $\square$

The following evaluations of q -binomial coefficients at a root of unity are well known, for instance see [13, Exercise 96 in Chapter 1].

Lemma 2.7. *Suppose w is a primitive d th root of unity and n is divisible by d . Then $[n]_{q=w} = n/d$ and*

$$\begin{bmatrix} n \\ k \end{bmatrix}_{q=w} = \begin{cases} \binom{n/d}{k/d} & \text{if } k \text{ is divisible by } d, \\ 0 & \text{otherwise.} \end{cases}$$

Now we are ready to prove Theorem 2.3.

Proof of Theorem 2.3. Suppose (c_1, c_2) is an (n, m) -annular rotation of order d . Then d divides both n and m . By Theorem 2.2 it is sufficient to show that for w a primitive d th root of unity, we have

$$(5) \quad X(w) = d \cdot \frac{(\widehat{n} - \widehat{R})(\widehat{m} - \widehat{S})}{\widehat{c}} \binom{\widehat{n}}{\widehat{r}} \binom{\widehat{m}}{\widehat{s}} \binom{\widehat{r}}{\widehat{\alpha}} \binom{\widehat{s}}{\widehat{\beta}} \binom{\widehat{c}}{\widehat{\lambda}} \binom{\widehat{c}}{\widehat{\mu}},$$

if all of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ are divisible by d , and $X(w) = 0$ otherwise. By Lemma 2.7 we get (5) when all of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ are divisible by d .

It remains to show that if at least one of $n, m, c, r, s, R, S, \alpha, \beta, \lambda, \mu$ is not divisible by d , then $X(w) = 0$. Since d divides both n and m , we have the following cases.

CASE 1: r or s is not divisible by d . By (4), $X(q)$ is a polynomial divisible by $\begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q$. Thus, by Lemma 2.7, we get $X(w) = 0$.

CASE 2: Both r and s are divisible by d , but α or β is not. Suppose $\alpha = (1^{a_1}, 2^{a_2}, \dots)$ is not divisible by d . Suppose moreover that a_j is not divisible by d . Again by (4), $X(q)$ is divisible by

$$\begin{bmatrix} r \\ \alpha \end{bmatrix}_q = \begin{bmatrix} r \\ a_j \end{bmatrix}_q \begin{bmatrix} r - a_j \\ a_1, \dots, a_{j-1}, a_{j+1}, \dots \end{bmatrix}_q.$$

Thus, by Lemma 2.7, we get $X(w) = 0$. If β is not divisible by d , by the same arguments we get $X(w) = 0$.

CASE 3: All of r, s, α, β are divisible by d , but c is not. Note that since α and β are divisible by d , so are R and S . By (4), we can write

$$X(q) = \frac{1 - q^{m-S}}{1 - q^c} Y(q)$$

for a polynomial $Y(q)$. Since $m - S$ is divisible by d , but c is not, $\frac{1 - w^{m-S}}{1 - w^c} = 0$ and we get $X(w) = 0$.

CASE 4: All of $r, s, \alpha, \beta, R, S, c$ are divisible by d , but λ or μ is not. Suppose λ is not divisible by d . By (4), $X(q)$ is divisible by $\begin{bmatrix} c \\ \lambda \end{bmatrix}_q$. By the same argument as in CASE 2 we obtain that $X(w) = 0$. If μ is not divisible by d , we can do the same thing using the following expression

$$X(q) = \frac{[(n-R)(m-S)]_q}{[n-R]_q} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} r \\ \alpha \end{bmatrix}_q \begin{bmatrix} s \\ \beta \end{bmatrix}_q \begin{bmatrix} c \\ \mu \end{bmatrix}_q \frac{[n-R]_q}{[c]_q} \begin{bmatrix} c \\ \lambda \end{bmatrix}_q.$$

Thus in all cases we have $X(w) = 0$, which finishes the proof. $\square$

3. ANNULAR q -KREWERAS NUMBERS

In this section we define annular versions of q -analogs of Kreweras, Narayana, and Catalan numbers and evaluate their sums.

For brevity we will use the following throughout this section:

$$X = c(c-1),$$

$$Y = r(c+r) + s(c+s),$$

$$Z = r(n-c-R) + s(m-c-S),$$

$$W = r(R-r) + s(S-s) + c(n-R-c) + c(m-S-c) - \tau(\alpha) - \tau(\beta) - \tau(\lambda) - \tau(\mu).$$

Definition 2. The *annular q -Kreweras number* $\text{Kre}^{\text{ann}} = \text{Kre}^{\text{ann}}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)$ is defined by

$$\text{Kre}^{\text{ann}} = q^X q^Y q^Z q^W \frac{[nm]_q}{[n]_q [m]_q} \frac{[2c]_q}{2} \frac{[n-R]_q [m-S]_q}{[c]_q^2} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} r \\ \alpha \end{bmatrix}_q \begin{bmatrix} s \\ \beta \end{bmatrix}_q \begin{bmatrix} c \\ \lambda \end{bmatrix}_q \begin{bmatrix} c \\ \mu \end{bmatrix}_q.$$

The *annular q -Narayana number* $\text{Nara}_1^{\text{ann}} = \text{Nara}_1^{\text{ann}}(n, m; c, r, s, R, S)$ of type 1 is defined by

$$\text{Nara}_1^{\text{ann}} = q^X q^Y q^Z \frac{[nm]_q}{[n]_q [m]_q} \frac{[2c]_q}{2} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} R-1 \\ r-1 \end{bmatrix}_q \begin{bmatrix} S-1 \\ s-1 \end{bmatrix}_q \begin{bmatrix} n-R \\ c \end{bmatrix}_q \begin{bmatrix} m-S \\ c \end{bmatrix}_q.$$

The *annular q -Narayana number* $\text{Nara}_2^{\text{ann}} = \text{Nara}_2^{\text{ann}}(n, m; c, r, s)$ of type 2 is defined by

$$\text{Nara}_2^{\text{ann}} = q^X q^Y \frac{[nm]_q}{[n]_q [m]_q} \frac{[2c]_q}{2} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} n \\ r+c \end{bmatrix}_q \begin{bmatrix} m \\ s+c \end{bmatrix}_q.$$

The *annular q -Narayana number* $\text{Nara}_3^{\text{ann}} = \text{Nara}_3^{\text{ann}}(n, m; , c)$ of type 3 is defined by

$$\text{Nara}_3^{\text{ann}} = q^X \frac{[nm]_q}{[n]_q [m]_q} \frac{[2c]_q}{2} \begin{bmatrix} 2n \\ n-c \end{bmatrix}_q \begin{bmatrix} 2m \\ m-c \end{bmatrix}_q.$$

The *annular q -Catalan number* $\text{Cat}^{\text{ann}}(n, m)$ is defined by

$$\text{Cat}^{\text{ann}}(n, m) = \frac{[nm]_q}{2[m+n]_q} \begin{bmatrix} 2n \\ n \end{bmatrix}_q \begin{bmatrix} 2m \\ m \end{bmatrix}_q.$$

In what follows we show that the above numbers are polynomials in q . Note however that they are not necessarily polynomials of integer coefficients. For instance, $\text{Cat}^{\text{ann}}(1, 1) = (1+q)/2$. We first show that Kre^{ann} is a polynomial in q .

Proposition 3.1. *The annular q -Kreweras number $\text{Kre}^{\text{ann}} = \text{Kre}^{\text{ann}}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)$ is a polynomial in q .*

Proof. Since

$$\text{Kre}^{\text{ann}} = q^X q^Y q^Z q^W \frac{[2c]_q}{2[c]_q} \left(\frac{[n-R]_q}{[n]_q} \begin{bmatrix} n \\ r \end{bmatrix}_q \begin{bmatrix} r \\ \alpha \end{bmatrix}_q \right) \left(\frac{[m-S]_q}{[c]_q} \begin{bmatrix} c \\ \mu \end{bmatrix}_q \right) \left(\frac{[nm]_q}{[m]_q} \begin{bmatrix} m \\ s \end{bmatrix}_q \begin{bmatrix} s \\ \beta \end{bmatrix}_q \begin{bmatrix} c \\ \lambda \end{bmatrix}_q \right),$$

we are done by Lemma 2.5. $\square$

In the introduction we saw that the sum of q -Kreweras numbers is equal to the q -Narayana number, and the sum of q -Narayana numbers is equal to the q -Catalan number. We show that annular versions of these numbers have similar properties. In order to do this we prove three lemmas.

The following lemma is due to Reiner and Sommers [10]. We include their elegant proof as well.

Lemma 3.2. [10] *Let $\tau(\lambda) = \sum_{i \geq 1} \lambda'_i \lambda'_{i+1}$, where λ' is the transposition of λ . Then*

$$\sum_{\lambda \in \text{Par}(n, k)} q^{k(n-k) - \tau(\lambda)} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q.$$


Proof. Let $\lambda = (1^{m_1}, 2^{m_2}, \dots)$. Then $m_1 + 2m_2 + \dots = n$ and $m_1 + m_2 + \dots = k$. Since $\lambda'_i = m_i + m_{i+1} + \dots$, we have

$$q^{k(n-k) - \tau(\lambda)} \begin{bmatrix} k \\ \lambda \end{bmatrix}_q = q^{(k-\lambda'_1)\lambda'_2} \begin{bmatrix} \lambda'_1 \\ \lambda'_2 \end{bmatrix}_q q^{(k-\lambda'_2)\lambda'_3} \begin{bmatrix} \lambda'_2 \\ \lambda'_3 \end{bmatrix}_q \dots.$$

Thus we can rewrite the identity as follows.

$$(6) \quad \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q = \sum_{\mu \in \text{Par}(n), \mu_1 = k} q^{(k-\mu_1)\mu_2} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}_q q^{(k-\mu_2)\mu_3} \begin{bmatrix} \mu_2 \\ \mu_3 \end{bmatrix}_q \dots.$$

The left hand side of (6) is the sum of $q^{|\nu|}$ for all partitions ν contained in a $k \times (n-k)$ rectangle where the k th part of ν is 0. For such a partition ν we define points $Q_1, P_1, Q_2, P_2, \dots$ as follows. Let Q_1 be the upper right corner of the rectangle. When Q_i is defined, let P_i be the point on the base of the rectangle which is straightly below Q_i . When P_i is defined, let Q_{i+1} be the intersection of the border of ν and the northwest diagonal ray starting from P_i . We define the sequence of points until we reach the bottom-left corner of the rectangle. Since ν has no cells in the k th row of the rectangle, we can always complete this sequence. Let μ be the partition whose i th part is equal to the length of the segment $P_{i-1}P_i$, where $P_0 = Q_1$. Then $\mu \in \text{Par}(n)$ and $\mu_1 = k$. It is easy to see that the sum of $q^{|\nu|}$ for partitions ν which give μ is equal to the summand in (6), see Figure 3. This proves (6). $\square$

FIGURE 3. We decompose ν and get μ .

Lemma 3.3. For fixed integers n, r, c , we have

$$\sum_{R \geq 0} q^{r(n-c-R)} \begin{bmatrix} R-1 \\ r-1 \end{bmatrix}_q \begin{bmatrix} n-R \\ c \end{bmatrix}_q = \begin{bmatrix} n \\ r+c \end{bmatrix}_q.$$

Proof. This can be proved by a standard technique considering the largest rectangle with width r contained in partitions inside an $(n-r-c) \times (r+c)$ rectangle. $\square$

Lemma 3.4. Let n, m , and k be any nonnegative integers. Then

$$\sum_{c \geq 0} q^{c(c-1+k)} [2c+k]_q \begin{bmatrix} 2n+k \\ n-c \end{bmatrix}_q \begin{bmatrix} 2m+k \\ m-c \end{bmatrix}_q = \frac{[n+k]_q [m+k]_q}{[n+m+k]_q} \begin{bmatrix} 2n+k \\ n+k \end{bmatrix}_q \begin{bmatrix} 2m+k \\ m+k \end{bmatrix}_q.$$

Proof. It is straightforward to check that

$$(7) \quad q^{c(c-1+k)} [n+m+k]_q [2c+k]_q = q^{c(c-1+k)} [n+c+k]_q [m+c+k]_q - q^{(c+1)(c+k)} [n-c]_q [m-c]_q.$$

Let $\sigma(c) = c(c-1+k)$. Then by (7) the left hand side is equal to

$$\begin{aligned} & \frac{1}{[n+m+k]_q} \sum_{c \geq 0} (q^{\sigma(c)} [n+c+k]_q [m+c+k]_q - q^{\sigma(c+1)} [n-c]_q [m-c]_q) \begin{bmatrix} 2n+k \\ n-c \end{bmatrix}_q \begin{bmatrix} 2m+k \\ m-c \end{bmatrix}_q \\ &= \frac{[2n+k]_q [2m+k]_q}{[n+m+k]_q} \sum_{c \geq 0} \left(q^{\sigma(c)} \begin{bmatrix} 2n+k-1 \\ n-c \end{bmatrix}_q \begin{bmatrix} 2m+k-1 \\ m-c \end{bmatrix}_q - q^{\sigma(c+1)} \begin{bmatrix} 2n+k-1 \\ n-c-1 \end{bmatrix}_q \begin{bmatrix} 2m+k-1 \\ m-c-1 \end{bmatrix}_q \right) \\ &= \frac{[2n+k]_q [2m+k]_q}{[n+m+k]_q} \begin{bmatrix} 2n+k-1 \\ n \end{bmatrix}_q \begin{bmatrix} 2m+k-1 \\ m \end{bmatrix}_q \\ &= \frac{[n+k]_q [m+k]_q}{[n+m+k]_q} \begin{bmatrix} 2n+k \\ n+k \end{bmatrix}_q \begin{bmatrix} 2m+k \\ m+k \end{bmatrix}_q. \end{aligned}$$

$\square$

Theorem 3.5. *We have*

$$\sum_{\substack{\alpha \in \text{Par}(R, r) \\ \beta \in \text{Par}(S, s) \\ \lambda \in \text{Par}(n-R, c) \\ \mu \in \text{Par}(m-S, c)}} \text{Kre}^{\text{ann}}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu) = \text{Nara}_1^{\text{ann}}(n, m; c, r, s, R, S),$$

$$\sum_{R, S \geq 0} \text{Nara}_1^{\text{ann}}(n, m; c, r, s, R, S) = \text{Nara}_2^{\text{ann}}(n, m; c, r, s),$$

$$\sum_{r, s \geq 0} \text{Nara}_2^{\text{ann}}(n, m; c, r, s) = \text{Nara}_3^{\text{ann}}(n, m; c),$$

$$\sum_{c \geq 0} \text{Nara}_3^{\text{ann}}(n, m; c) = \text{Cat}^{\text{ann}}(n, m).$$

Proof. The first, second, and fourth identities follow from Lemmas 3.2, 3.3, and 3.4, respectively. The third identity follows from the q -Vandermonde's identity:

$$\sum_{i \geq 0} q^{i(m-k+i)} \begin{bmatrix} m \\ k-i \end{bmatrix}_q \begin{bmatrix} n \\ i \end{bmatrix}_q = \begin{bmatrix} m+n \\ k \end{bmatrix}_q.$$

□

It is easy to see that we can replace the polynomial $X(q)$ in Theorem 2.3 with Kre^{ann} . Thus by Theorems 2.3 and 3.5 we obtain the following.

Theorem 3.6. *The following exhibit the cyclic sieving phenomenon:*

$$\begin{aligned} &(\text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu), C, \text{Kre}^{\text{ann}}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu)), \\ &(\text{ANC}(n, m; c, r, s, R, S), C, \text{Nara}_1^{\text{ann}}(n, m; c, r, s, R, S)), \\ &(\text{ANC}(n, m; c, r, s), C, \text{Nara}_2^{\text{ann}}(n, m; c, r, s)), \\ &(\text{ANC}(n, m; c), C, \text{Nara}_3^{\text{ann}}(n, m; c)), \\ &(\text{ANC}(n, m), C, \text{Cat}^{\text{ann}}(n, m)). \end{aligned}$$

Considering the annular rotation of order 1 in Theorem 3.6, i.e. the identity action, we obtain the following.

Corollary 3.7. *We have*

$$(8) \quad \# \text{ANC}(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu) = \frac{(n-R)(m-S)}{c} \binom{n}{r} \binom{m}{s} \binom{r}{\alpha} \binom{s}{\beta} \binom{c}{\lambda} \binom{c}{\mu},$$

$$(9) \quad \# \text{ANC}(n, m; c, r, s, R, S) = c \binom{n}{r} \binom{m}{s} \binom{R-1}{r-1} \binom{S-1}{s-1} \binom{n-R}{c} \binom{m-S}{c},$$

$$(10) \quad \# \text{ANC}(n, m; c, r, s) = c \binom{n}{r} \binom{m}{s} \binom{n}{r+c} \binom{m}{s+c},$$

$$(11) \quad \# \text{ANC}(n, m; c) = c \binom{2n}{n-c} \binom{2m}{m-c},$$

$$(12) \quad \# \text{ANC}(n, m) = \frac{nm}{2(m+n)} \binom{2n}{n} \binom{2m}{m}.$$

If an annular noncrossing permutation is invariant under an annular rotation of order 2, it is called an *annular noncrossing permutation of type B*. We define $\text{ANC}_B(n, m)$ to be the set of connected $(2n, 2m)$ -annular noncrossing permutations of type B . We then define

$$\text{ANC}_B(n, m; c) = \text{ANC}_B(n, m) \cap \text{ANC}(2n, 2m; 2c),$$

$$\text{ANC}_B(n, m; c, r, s) = \text{ANC}_B(n, m) \cap \text{ANC}(2n, 2m; 2c, 2r, 2s),$$

$$\text{ANC}_B(n, m; c, r, s, R, S) = \text{ANC}_B(n, m) \cap \text{ANC}(2n, 2m; 2c, 2r, 2s, 2R, 2S),$$

$$\text{ANC}_B(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu) = \text{ANC}_B(n, m) \cap \text{ANC}(2n, 2m; 2c, 2r, 2s, 2R, 2S; 2\alpha, 2\beta, 2\lambda, 2\mu),$$

where $2\lambda = (2\lambda_1, 2\lambda_2, \dots)$.

Notice that every connected annular noncrossing permutation of type B contains at least two connected cycles. Thus annular noncrossing permutations of type B are in bijection with annular noncrossing partitions of type B .

Considering an annular rotation of order 2 in Theorem 3.6 we obtain the following.

Corollary 3.8. *We have*

$$(13) \quad \# \text{ANC}_B(n, m; c, r, s, R, S; \alpha, \beta, \lambda, \mu) = \frac{2(n-R)(m-S)}{c} \binom{n}{r} \binom{m}{s} \binom{r}{\alpha} \binom{s}{\beta} \binom{c}{\lambda} \binom{c}{\mu},$$

$$(14) \quad \# \text{ANC}_B(n, m; c, r, s, R, S) = 2c \binom{n}{r} \binom{m}{s} \binom{R-1}{r-1} \binom{S-1}{s-1} \binom{n-R}{c} \binom{m-S}{c},$$

$$(15) \quad \# \text{ANC}_B(n, m; c, r, s) = 2c \binom{n}{r} \binom{m}{s} \binom{n}{r+c} \binom{m}{s+c},$$

$$(16) \quad \# \text{ANC}_B(n, m; c) = 2c \binom{2n}{n-c} \binom{2m}{m-c},$$

$$(17) \quad \# \text{ANC}_B(n, m) = \frac{nm}{m+n} \binom{2n}{n} \binom{2m}{m}.$$

We note that (12) was first proved by Mingo and Nica [8, Corollary 6.7] and (15), (16), and (17) were first proved by Goulden, Nica and Oancea [4, Equations (4.6), (4.7), (4.9)].

4. ANNULAR NONCROSSING MATCHINGS

An (n, m) -annular noncrossing matching is a complete matching on $[n + m]$ which can be drawn in an (n, m) -annulus without crossing. By considering each matching pair (i, j) as a cycle of size 2, one can identify an annular noncrossing matching as an annular noncrossing permutation consisting of cycles of size 2 only.

Theorem 4.1. *Suppose $n \equiv m \equiv c \pmod{2}$. The number of (n, m) -annular noncrossing matchings exactly c connected matching pairs is*

$$c \binom{n}{\frac{n-c}{2}} \binom{m}{\frac{m-c}{2}}.$$

Proof. Such an annular noncrossing matching can be considered as an annular noncrossing permutation with $\alpha = (2^R) \in \text{Par}(2R, R)$, $\beta = (2^S) \in \text{Par}(2S, S)$, $\lambda = \mu = (1^c) \in \text{Par}(c, c)$, and $n - 2R = m - 2S = c$. Then we obtain the formula immediately from (8). $\square$

We can also obtain the total number of connected annular noncrossing matchings.

Theorem 4.2. *For $n \equiv m \pmod{2}$, the number of connected (n, m) -annular noncrossing matchings is*

$$\frac{2 \lceil \frac{n}{2} \rceil \lceil \frac{m}{2} \rceil}{n + m} \binom{n}{\lceil \frac{n}{2} \rceil} \binom{m}{\lceil \frac{m}{2} \rceil}.$$

Proof. We will prove the equivalent statement: for $k \in \{0, 1\}$, the number of connected $(2n + k, 2m + k)$ -annular noncrossing matchings is

$$\frac{(n+k)(m+k)}{n+m+k} \binom{2n+k}{n+k} \binom{2m+k}{m+k}.$$

By Theorem 4.1 the number is equal to

$$\sum_{c \geq 0} (2c + k) \binom{2n+k}{n-c} \binom{2m+k}{m-c}.$$

Then we are done by Lemma 3.4 for $q = 1$. $\square$

ACKNOWLEDGEMENT

The author would like to thank Vic Reiner for providing his paper with Sommers [10], suggesting the problem of finding a bicyclic sieving phenomenon on annular noncrossing permutations, and for fruitful comments. He also thanks Dennis Stanton for helpful discussion.

REFERENCES

- [1] H. Barcelo, V. Reiner, and D. Stanton. Bimahonian distributions. *J. Lond. Math. Soc. (2)*, 77(3):627–646, 2008.
- [2] D. Bessis and V. Reiner. Cyclic sieving of noncrossing partitions for complex reflection groups. *Ann. Comb.*, 15(2):197–222, 2011.
- [3] I. P. Goulden and D. M. Jackson. Transitive factorizations into transpositions and holomorphic mappings of the sphere. *Proc. Amer. Math. Soc.*, 125:51–60, 1997.
- [4] I. P. Goulden, A. Nica, and I. Oancea. Enumerative properties of $NC^{(B)}(p, q)$. *Ann. Comb.*, 15(2):277–303, 2011.
- [5] J. S. Kim. Chain enumeration of k -divisible noncrossing partitions of classical types. *J. Combin. Theory Ser. A*, 118(3):879–898, 2011.
- [6] J. S. Kim, S. Seo, and H. Shin. Annular noncrossing permutations and minimal transitive factorizations. <http://arxiv.org/abs/1201.5703>.
- [7] G. Kreweras. Sur les partitions non croisées d’un cycle. *Discrete Math.*, 1:333–350, 1972.
- [8] J. A. Mingo and A. Nica. Annular noncrossing permutations and partitions, and second-order asymptotics for random matrices. *Int. Math. Res. Not.*, (28):1413–1460, 2004.
- [9] A. Nica and I. Oancea. Posets of annular non-crossing partitions of types B and D . *Discrete Math.*, 309(6):1443–1466, 2009.
- [10] V. Reiner and E. Sommers. q -analogues of Weyl group Narayana and Kreweras numbers. preprint.
- [11] V. Reiner, D. Stanton, and D. White. The cyclic sieving phenomenon. *J. Combin. Theory Ser. A*, 108(1):17–50, 2004.
- [12] B. E. Sagan. *The cyclic sieving phenomenon: a survey*, volume 392 of *London Math. Soc. Lecture Note Ser.* Cambridge Univ. Press, Cambridge, 2011.
- [13] R. P. Stanley. *Enumerative Combinatorics. Vol. 1, second ed.* Cambridge University Press, New York/Cambridge, 2011.